

AD-A117 085

YALE UNIV NEW HAVEN CT COWLES FOUNDATION FOR RESEARC--ETC F/8 12/1  
STOCHASTIC GAMES. I. FOUNDATIONS.(U)  
APR 82 C A MONASH  
DISCUSSION PAPER-623

N00014-77-C-0518

NL

UNCLASSIFIED

[ ]  
AL  
DATE



END  
DATE  
FILED  
82  
DTIC

AD A117085

COWLES FOUNDATION FOR RESEARCH IN ECONOMICS

AT YALE UNIVERSITY

Box 2125, Yale Station  
New Haven, Connecticut 06520

COWLES FOUNDATION DISCUSSION PAPER NO.-623

*Contract N00014-77-C-0518*

DTIC  
ELECTE  
JUL 20 1982

Note: Cowles Foundation Discussion Papers are preliminary materials circulated to stimulate discussion and critical comment. Requests for single copies of a Paper will be filled by the Cowles Foundation within the limits of the supply. References in publications to Discussion Papers (other than mere acknowledgment by a writer that he has access to such unpublished material) should be cleared with the author to protect the tentative character of these papers.

STOCHASTIC GAMES I: Foundations

by

Curt Alfred Monash

April 1982

DTIC FILE COPY

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

①

STOCHASTIC GAMES I: FOUNDATIONS

by

Curt Alfred Monash

DTIC  
JUL 29 1982

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution unlimited

1.

This paper strives to provide a sound underpinning for the theory of stochastic games. Section 2 is a reworking of the Bewley-Kohlberg result integrated with Shapley's; the "black magic" of Tarski's principle is replaced by the "gray magic" of the Hilbert Nullstellensatz. Section 3 explicates the underlying topology and measure theory; I believe it is as necessary for Mertens and Neyman's proof of the minmax theorem [8] as it is for mine [9, 10]. Finally, Section 4 establishes a result on this sort of structure which may be of some independent interest; in any case, it is critical for the argument in [10].

This work was strongly influenced by Andrew Gleason. Other acknowledgments may be found in [10].

Accession For	
NTIS	<input checked="checked" type="checkbox"/>
ERIC	<input type="checkbox"/>
Other	<input type="checkbox"/>
Per	<input type="checkbox"/>
FL-171	on file
A	

2.

Fix a stochastic game. In this section we study the r-discount game, following [11] and [3].

Let  $S$ , the state set,  $A$  (resp.  $B$ ) the choice set each turn for  $A$  (resp.  $B$ ), and  $C$ , with fixed measurable function  $q: [0,1] \rightarrow C$  all be finite sets; let  $\delta: S \times A \times B \times C \rightarrow S$  and  $d: S \times A \times B \times C \rightarrow [-\hat{M}, \hat{M}]$  give the state and the outcome resulting from choices (starting in state  $s \in S$ ) of  $a \in A$ ,  $b \in B$  and random  $c \in C$ . Without loss of generality, these sets and functions describe the game completely. Consider now the stochastic game, with fixed starting state  $s$ , and discount rate  $r \in (0,1)$ ; that is, let the payoff function be  $\sum_{i=1}^{\infty} d_i(1-r)^{i-1}$ , where  $d_i$  the payoff on the  $i^{\text{th}}$  step.

As  $\sum_{i=1}^{\infty} d_i(1-r)^{i-1}$  is a continuous payoff function on a pair of compact strategy spaces (see Section 3) the min-max theorem follows immediately (this well-known result follows rapidly from [11]); let the value of the game, for starting state  $s$ , be  $V_s(r)$ . Following [3] and [11] we characterize the values and optimal strategies, by considering the system of equations:

$$U_s(r) = \text{val}_C(\text{Exp}(d(s,a,b,c)) + (1-r) \sum_{s \in S} P(\bar{s}) \cdot U_s(r)) . \quad (2.1s)$$

Here  $P(\bar{s}) = \text{measure}(\{t \in [0,1] : \delta(s,a,b,q(t)) = \bar{s}\})$  depends upon  $a$ ,  $b$ , and  $s$ ;  $\text{val}$  is just the ordinary min-max value of the matrix. For fixed  $r$ , think of the system as one equation in a variable in  $c^{|S|}$ . Applying the Contraction Mapping Theorem ([6], p. 229), it must have a unique solution  $(\hat{U}_s(r))$ . Indeed, as the contraction constant  $\|1-r\|$  is bounded away from 1 on any interval  $[r^*, 1]$ , and indeed on a set  $\{x+y \in C : r^* \leq x \leq 1, 0 \leq y \leq \delta(r^*)\}$ ,  $(\hat{U}_s(r))$  is

a continuous function of  $r$  in (a neighborhood of)  $(0,1)$  .

Proposition 2.2: For  $r \in (0,1)$  , this solution is precisely  $(V_s(r))$  .

Proof: It is clearly enough to show the following, for fixed  $r \in (0,1)$  :

By playing as his  $i^{\text{th}}$  move an optimal strategy in the game given by the matrix of  $(2.1s_{i-1})$ , Mr. A (2.3) guarantees himself a payoff of at least  $\tilde{U}_{s_0}(r)$  .

(2.3) is clearly equivalent to:

$$\tilde{U}_{s_0}(r) \geq V_{s_0}(r) \text{ for all } s_0 \in S . \quad (2.4)$$

So suppose (2.4) is false. Let  $\alpha = \max_S (V_{s_0}(r) - \tilde{U}_{s_0}(r))$  ; choose  $\tilde{s}$  so that the max is achieved.

Then

$$\begin{aligned} \tilde{U}_{\tilde{s}}(r) &= \text{val}_C(\text{Exp}(d(\tilde{s}, a, b, c))) + \sum_{s \in S} (1-r)P(\bar{s}) \cdot \tilde{U}_{\bar{s}}(r) \\ &= -(1-r)\alpha + \text{val}_C(\text{Exp}(d(\tilde{s}, a, b, c))) + \sum_{s \in S} (1-r)P(\bar{s}) \cdot (\tilde{U}_{\bar{s}}(r) + \alpha) \\ &\geq -(1-r)\alpha + \text{val}_C(\text{Exp}(d(\tilde{s}, a, b, c))) + \sum_{s \in S} (1-r)P(s) \cdot V_{\bar{s}}(r) . \end{aligned}$$

$$\text{So } \tilde{U}_{\tilde{s}}(r) + \alpha > \text{val}_C(\text{Exp}(d(\tilde{s}, a, b, c))) + \sum_{s \in S} (1-r)P(s; a, b, \tilde{s}) \cdot V_{\bar{s}}(r) . \quad (2.5)$$

But since both sides of (2.5) equal  $V_{\tilde{s}}(r)$  , we have a contradiction.

□

It is well-known that the value of a matrix is one of a finite list of rational functions of the coefficients (namely the value of some square

submatrix; see [8], p. 76). Thus the system may be viewed as a set of polynomial equations--identifying  $S$  as  $\{1, \dots, k\}$ , where  $k = |S|$ --  
 $g_1(x_1, \dots, x_k) = 0, \dots, g_k(x_1, \dots, x_k) = 0$ , where the coefficients  
 lie in the algebraically closed field  $F$ . Now, if the ideal

$(g_1, \dots, g_k)$  --that is,  $\{\sum_{i=1}^k h_i g_i : h_i \in F, i = 1, \dots, k\}$  --equals the  
 unit ideal  $(1)$ , then some linear combination  $\sum_{i=1}^k h_i g_i = 1, h_i \in F$   
 for  $i = 1, \dots, k$ ; this is impossible because  $\sum_{i=1}^k h_i g_i((V_S)) = \sum_{i=1}^k h_i \cdot 0 \equiv 0$   
when viewed as functions of  $r$ .

By the Hilbert Nullstellensatz (weak form; see [2], p. 69), then,  
 the only other possibility is that  $\{(x_1, \dots, x_k) \in F^k : g_1(x_1, \dots, x_k) = 0, \dots, g_k(x_1, \dots, x_k) = 0\}$  is non-empty. On the other hand, we saw  
 above that the solution is unique in the much larger field of all functions  
 of  $r$ ; hence it is certainly unique in  $F$ , and precisely equal to  
 $(V_S(r))$ . But an algebraic function which maps reals to reals must in-  
 deed lie in  $F' =$  the field of real algebraic functions  $\subset F$ . Finally,  
 observe that each  $V_S(r)$  equals, at each point, one of a finite list of  
 algebraic functions; each pair of these can cross each other only finitely  
 often (for their difference is an algebraic function, which can only have  
 finitely many zeroes). Thus by the continuity of the  $V_S(r)$ , there  
 exists an interval  $(0, \tilde{r})$  on which the  $V_S(r)$  are algebraic, for all  
 $s \in S$ .

$F'$  has a natural ordering:  $f > g$  if and only if there exists  
 an  $\tilde{r}$  such that  $f > g$  on the interval  $(0, \tilde{r})$ . Consider the games  
 (2.1s) with the values  $V_S(r)$  substituted in place of the  $U_S(r)$ ; then  
 these are ordinary two-person zero-sum games, with coefficients in  $F'$ ,  
 and hence have optimal strategies  $\{f(s, s; r)\}$ ; by construction, for  
 each  $s$  and  $s$ ,  $f(s, s; r)$  belongs to  $F'$ . For each  $r$ , then, we

have optimal stationary strategies, algebraic in  $r$ , for the  $r$ -discount game. Finally, it is well known that algebraic functions have Puiseux (fractional power) series expansions—see [1]—and so we have checked the

Theorem of Bewley and Kohlberg: The values of the  $r$ -discount game, and the stationary optimal strategies, have Puiseux expansions.



3.

More generally, consider an infinite decision tree such that, at each node, the choice set is finite. By possibly duplicating certain branches, we may assume that, without loss of generality, there exist finite sets  $Z_i$ ,  $i = 1, 2, \dots$ , such that  $Z_1$  consists simply of the first node,  $Z_2$  is the choice set at the first node,  $Z_3$  is the choice set at the second node, no matter which node was chosen as the second node, and so on. Let  $Q$  be the space of paths on the tree;

$Q = Z_1 \times Z_2 \times \dots$  is compact by Tychonoff's theorem ([9], p. 144).

Simultaneously fix probability distributions on the choice set at each node. Then these distributions induce a sequence of functions  $\mu_i : Z_1 \times \dots \times Z_i \rightarrow \mathbb{R}$  such that

$$\sum_{z \in Z_i} \mu_i(z_1, z_2, \dots, z_{i-1}, z) = \mu_{i-1}(z_1, \dots, z_{i-1})$$

for any fixed  $z_1, \dots, z_{i-1}$ , for any  $i = 1, 2, \dots$ .

The following is a special case of the Kolmogorov Extension Theorem ([5]).

Proposition 3.1: There exists a measure  $\mu$  on  $Q$  such that

$\mu(\{z_1\} \times \{z_2\} \times \dots \times \{z_i\} \times Z_{i+1} \times Z_{i+2} \times \dots) = \mu_i(z_1, \dots, z_i)$ , for all  $z_1, \dots, z_i$ , for all  $i$ .

Proof: Consider the ring generated by  $\mathcal{Z}$  = the collection of sets of the form

$$\{z_1\} \times \dots \times \{z_i\} \times Z_{i+1} \times Z_{i+2} \times \dots$$

We clearly have a finitely additive set function  $\mu$  with the desired

properties; to check that  $\mu$  is a measure, we need only confirm that for any descending tower  $Y_1 \supseteq Y_2 \supseteq Y_3 \supseteq \dots$ ,  $Y_j \in \mathcal{Z}$  for all  $j$ ,  $\bigcap_{j=1}^{\infty} Y_j = \phi$  implies  $\lim_{j \rightarrow \infty} \mu(Y_j) = 0$  ([4], p. 39). But every element of  $\mathcal{Z}$  is both open and closed in the direct product topology; hence the same is true of every member of the ring. Thus, by the compactness of  $Q$ , any tower descending to 0 has  $Y_j = \phi$  for all  $j$  sufficiently large; hence  $\lim_{j \rightarrow \infty} \mu(Y_j) = 0$  trivially. Finally, since  $\mathcal{Z}$  is a base for the direct product topology, the Caratheodory Extension Theorem extends to the Borel field ([4], p. 54).



Denote by  $\tau$  a function which assigns to every node in  $N$ , the set of nodes, a probability distribution on its choice set; denote the set of all possible  $\tau$  by  $M$ , the space of mixed strategies. Then every  $\tau \in M$  induces a probability measure  $\mu_\tau$  on  $Q$ . Let  $P \in M$ , the space of pure strategies, be the set of all  $\pi \in M$  such that, at every node,  $\pi$  selects some alternative with probability one. Then there exists a canonical mapping  $f: P \rightarrow Q$ , sending  $\pi \in P$  into the path determined by  $\pi$ .  $\tau$  also induces a probability measure  $\theta_\tau$  on  $P$ , and we have  $\theta_\tau = \mu_\tau \circ f$ .

Partition  $N$  into two disjoint subsets  $N_B \cup N_C$  where  $N_B$  will be thought of as the set of decision nodes, and  $N_C$  as the set of chance nodes. Then we can write  $M = M_B \times M_C$ , where an element  $\tau_B(\tau_C)$  of  $M_B(M_C)$  is an assignment to each node in  $N_B(N_C)$  a probability distribution on its choice set. Similarly  $P = P_B \times P_C$ .

$\theta(\tau_B, \tau_C)$  then decomposes as a product measure:

$$\theta_{(\tau_B, \tau_C)} = \theta_{\tau_B} \times \theta_{\tau_C}.$$

Assume now that  $\tau_C$  is fixed. Letting  $d_i$ ,  $i = 1, 2, \dots$  be a sequence of measurable functions on  $Q$ , bounded below, we have

Proposition 3.2:

$$\begin{aligned} & \inf_{\tau_B \in M_B} \liminf_{N \rightarrow \infty} \int_Q \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\mu_{(\tau_B, \tau_C)} \\ &= \inf_{\pi_B \in P_B} \liminf_{N \rightarrow \infty} \int_Q \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\mu_{(\pi_B, \pi_C)}. \end{aligned}$$

Thus thinking of  $N_B$  as the nodes controlled by the decision-maker and  $N_C$  as those controlled by nature (with nature's "strategy" known in advance, the decision-maker "might as well" play a pure strategy as a mixed one. This is true even in a slightly stronger sense:

Corollary 3.3: If there exists a mixed strategy such that the infimum is achieved, then it is also achieved by a pure strategy.

Proof of Proposition 3.2: For any  $i$ , for any  $\tau_B$ ,

$$\begin{aligned} \int_Q d_i d\mu_{(\tau_B, \tau_C)} &= \int_P d_i(f(\pi)) d\theta_{(\tau_B, \tau_C)}(\pi) \\ &= \int_{P_B} \left( \int_{P_C} d_i(f(\pi_B, \pi_C)) d\theta_{\tau_B}(\pi_C) \right) d\theta_{\tau_B}(\pi_B) \end{aligned}$$

by Fubini's Theorem ([9], p. 233). Thus of course

$$\int_Q \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\mu_{(\tau_B, \tau_C)} = \int_{P_B} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} d\theta_{\tau_B}.$$

Observing that the support of  $\theta_{\pi_B}$  is a subset of  $P_B$  on which  $f(\pi, \pi_C) \equiv f(\pi_B, \pi_C)$ , for any  $\pi_C \in P_C$ , we see:

$$\begin{aligned} & \inf_{\pi_B \in P_B} \liminf_{N \rightarrow \infty} \int_{P_B} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} d\theta_{\tau_B} \\ &= \inf_{\pi_B \in P_B} \liminf_{N \rightarrow \infty} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C}. \end{aligned}$$

But for any fixed  $\tau_B$ ,

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \int_{P_B} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} d\theta_{\tau_B} \\ & \geq \inf_{\pi_B \in P_B} \liminf_{N \rightarrow \infty} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C}, \end{aligned}$$

by Fatou's Lemma ([9], p. 199); of course, the same inequality holds if we replace the left-hand side by the  $\inf_{\tau_B \in M_B}$ . But, since  $P_B \subset M_B$ , the reverse inequality also holds and so

$$\begin{aligned} & \inf_{\tau_B \in M_B} \liminf_{N \rightarrow \infty} \int_{P_B} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} d\theta_{\tau_B} \\ &= \inf_{\pi_B \in P_B} \liminf_{N \rightarrow \infty} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C}, \text{ and done.} \end{aligned}$$

□

Proof of Corollary 3.3: For  $\tau_B$  such that the infimum is achieved, the

set of  $\pi$  in the support of  $\theta_{\tau_B}$  such that  $\liminf_{N \rightarrow \infty} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} \geq I + \frac{1}{k}$  is clearly of measure zero, where  $I$  is the infimum, and  $k$  an arbitrary

positive integer. So the set where

$$\liminf_{N \rightarrow \infty} \int_{P_C} \left( \frac{1}{N} \sum_{i=1}^N d_i \right) d\theta_{\tau_C} \neq 1$$

is indeed of measure zero. Since the support of  $\theta_{\tau_B}$  is of course of measure one, we are done.



Note 3.4: The results of this section remain valid if "finite" is replaced by "compact."

Note 3.5: Our "mixed strategies" may not be the most general the decision-maker could adopt. However, strategies with more complicated forms of mixing can be disposed of in the same fashion using the compact form of Proposition 3.2 (see Note 3.4).

4.

Let  $\{Z_i\}$ ,  $Q$ ,  $\{\nu_i\}$ , and  $\nu$  be as in the previous section.

Suppose each  $Z_i$  has a distinguished subset  $Z_i^*$  such that

$(Z_1 \times \dots \times Z_{i-1} \times Z_i^* \times (Z_{i+1}^*)^c \times Z_{i+2}^* \times \dots) = \emptyset$ ; in other words, once a node in some  $Z_i^*$  is reached, the path remains in the  $\{Z_i^*\}$  with probability 1. For a given path  $z = (z_1, z_2, \dots)$ , let

$$P_*(i; z) = \begin{cases} \frac{\nu(\{z_1\} \times \dots \times \{z_{i-1}\} \times Z_i^* \times Z_{i+1} \times \dots)}{\nu(\{z_1\} \times \dots \times \{z_{i-1}\} \times Z_i \times Z_{i+1} \times \dots)} & \text{if } z_{i-1} \notin Z_{i-1}^* \\ 0 & \text{if } z_i \in Z_{i-1}^* \end{cases}$$

the probability (conditional on  $z_1, \dots, z_{i-1}$ ) of entering  $Z_i^*$ . Suppose

we wish to integrate a function over  $Q$  which vanishes outside

$(Z_1^*)^c \times (Z_2^*)^c \times \dots$ . Then, intuitively, we can integrate a smaller function, on a new measure which is bigger on  $(Z_1^*)^c \times (Z_2^*)^c \times \dots$ , to achieve the same result. This can be helpful if we wish to estimate the integral by estimating the (smaller) function directly.

More precisely, we prove this result in terms of truncated sequences;

it is applied in Section 5 of [10]. For any  $z = (z_1, z_2, \dots) \in Q$ ,

define the truncations  $z(k) = (z_1, \dots, z_k) \in Z_1 \times \dots \times Z_k$ . Write also

$Q(z_1, \dots, z_k) = \{z_1\} \times \{z_2\} \times \dots \times \{z_k\} \times Z_{k+1} \times Z_{k+2} \times \dots \subset Q$ . Then

**Proposition 4.1:** There exists a probability measure  $\tilde{\nu}$  on  $Q$  such that,

for all  $k$ , for all  $f_k : Z_1 \times \dots \times Z_k \rightarrow \mathbb{R}$  such that

$f_k : Z_1 \times \dots \times Z_{k-1} \times Z_k^* \rightarrow \{0\}$ ,

$$\int_Q f_k(z(k)) d\nu = \int_Q f_k(z(k)) \cdot \prod_{i=1}^k (1 - P_*(i; z)) d\tilde{\nu}.$$

Proof: By Proposition 2.1, to specify  $\tilde{\mu}$  we need only specify  $\tilde{\mu}(Q(z_1, \dots, z_i))$ , for all  $z_1, \dots, z_i$ , for all  $i$ , while checking that

$$\sum_{\bar{z}_i \in Z_i} \tilde{\mu}(Q(z_1, \dots, z_{i-1}, \bar{z}_i)) = \tilde{\mu}(Q(z_1, \dots, z_{i-1})) \quad (4.2)$$

So we define  $\tilde{\mu}(Q(z_1, \dots, z_k))$  inductively on  $k$ .

$\tilde{\mu}(Q) = 1$ . Fix  $z = (z_1, \dots) \in Q$

$$\tilde{\mu}(Q(z_1)) = \begin{cases} \frac{1}{1 - P_*(1; z)} \cdot \mu(Q(z_1)) & \text{if } z_1 \notin Z_1^* \\ 0 & \text{if } z_1 \in Z_1^* \end{cases} \quad (4.3)$$

unless  $P_*(1; z) = 1$ . In that case,

$$\tilde{\mu}(Q(z_1)) = \mu(Q(z_1)).$$

More generally, suppose we have specified  $\tilde{\mu}(Q(z_1, \dots, z_{i-1}))$ . If this equals 0, we let  $\tilde{\mu}(Q(z_1, \dots, z_{i-1}, z_i)) = 0$ . If not, then also  $\mu(Q(z_1, \dots, z_{i-1})) \neq 0$ , and we let

$$\tilde{\mu}(Q(z_1, \dots, z_i)) = \begin{cases} \frac{1}{1 - P_*(i; z)} \cdot \frac{\tilde{\mu}(Q(z_1, \dots, z_{i-1}))}{\mu(Q(z_1, \dots, z_{i-1}))} \cdot \mu(Q(z_1, \dots, z_i)) & \text{if } z_i \notin Z_i^* \\ 0 & \text{if } z_i \in Z_i^* \end{cases}$$

unless, again,  $P_*(i; z) = 1$ , in which case we let  $\tilde{\mu}(Q(z_1, \dots, z_i)) = \mu(Q(z_1, \dots, z_i))$ .

Now

$$\sum_{\bar{z}_1 \in Z_1} \tilde{\mu}(z_1, \dots, z_{i-1}, \bar{z}_1) \\ = \frac{1}{1 - P_*(i; z)} \frac{\tilde{\mu}(Q(z_1, \dots, z_{i-1}))}{\mu(Q(z_1, \dots, z_{i-1}))} \sum_{\bar{z}_1 \notin Z_1^*} \mu(Q(z_1, \dots, z_{i-1}, \bar{z}_1)) .$$

But 
$$\sum_{\bar{z}_1 \notin Z_1^*} \mu(Q(z_1, \dots, z_{i-1}, \bar{z}_1)) = (1 - P_*(i; z)) \mu(Q(z_1, \dots, z_{i-1}))$$

by the definition of  $P_*(i; z)$ , and so (4.2) is checked, and we have defined a measure  $\tilde{\mu}$ .

It remains to show that

$$\int_Q f_k(z(k)) d\mu = \int_Q f_k(z(k)) \cdot \prod_{i=1}^k (1 - P_*(i; z)) d\tilde{\mu} .$$

We start by observing that

$$\int_Q = \sum_{z_1 \times \dots \times z_k} \int_{Q(z_1 \times \dots \times z_k)} .$$

But on  $Q(z_1 \times \dots \times z_k)$ ,  $f_k$  is just a constant, as is  $\prod_{i=1}^k (1 - P_*(i; z))$ ,

and hence we are asking whether  $\int f(z_1, \dots, z_k) \cdot \tilde{\mu}(Q(z_1, \dots, z_k))$

$$= \int f(z_1, \dots, z_k) \cdot \prod_{i=1}^k (1 - P_*(i; z)) \tilde{\mu}(Q(z_1, \dots, z_k)) .$$

This of course follows

if

$$\mu(Q(z_1, \dots, z_k)) = \prod_{i=1}^k (1 - P_*(i; z)) \cdot \tilde{\mu}(z_1, \dots, z_k) \quad (4.4)$$

whenever  $f_k(z_1, \dots, z_k) \neq 0$  —i.e., whenever  $z_k \notin Z_k^*$ .

When  $k = 1$ , (4.4) is immediate by (4.3). So assume proved for  $k-1$ , and check for  $k$ . Then, for any  $z_1 \notin Z_1^*$ , we define  $\bar{\mu}$  on



$Q(z_1)$  by  $\bar{\mu} \equiv \frac{1}{1 - P_*(1; z)} \cdot \mu$ . Defining  $\tilde{\mu}$  exactly as before, we apply induction to get

$$\begin{aligned} & \prod_{i=1}^k (1 - P_*(1; z)) \cdot \tilde{\mu}(Q(z_1, \dots, z_k)) \\ &= (1 - P_*(1; z)) \cdot \bar{\mu}(Q(z_1, \dots, z_k)) = \mu(Q(z_1, \dots, z_k)) , \end{aligned}$$

and we are done.



## BIBLIOGRAPHY

- [1] Ahlfors, L. (1966). Complex Analysis, second edition. New York: McGraw-Hill.
- [2] Atiyah, M. F. and I. G. McDonald (1969). Introduction to Commutative Algebra. Reading, Mass.: Addison-Wesley.
- [3] Bewley, T. and E. Kohlberg (1976). "The Asymptotic Theory of Stochastic Games," Math. Oper. Res., 1, 197-208.
- [4] Halmos, P. (1950). Measure Theory. New York: Van Nostrand.
- [5] Kolmogorov, A. N. (1950). Foundations of the Theory of Probabilities, translated by Nathan Morrison. New York: Chelsea Publishing Co.
- [6] Loomis, L. and S. Sternberg (1968). Advanced Calculus. Reading, Mass.: Addison-Wesley.
- [7] Mertens, J. F. and A. Neyman (1980). "Stochastic Games," C.O.R.E. Report No. 22/80.
- [8] McKinsey, J. (1952). Introduction to the Theory of Games. New York: McGraw-Hill.
- [9] Monash, C. A. (1979). "Stochastic Games: the Minmax Theorem," Ph.D Dissertation, Mathematics Department, Harvard University.
- [10] \_\_\_\_\_ (1981). "Stochastic Games II: The Minmax Theorem," Cowles Foundation Discussion Paper 624.
- [11] Shapley, L. (1953). "Stochastic Games," Proc. Nat. Acad. Sci. (USA), 39, 1095-1100.
- [12] Wald, A. (1945). "Generalization of a Theorem by von Neumann Concerning Zero Sum Two Person Games," Annals of Mathematics, 46, 281-286.

# ACKNOWLEDGMENTS

Partially supported by the Office of Naval Research Contract Number  
N00014-77-0518 and a National Science Foundation Graduate Fellowship.